

BIBLIOGRAFÍA RECOMENDADA EN BIOPSIA SELECTIVA DE GANGLIO CENTINELA (BSGC)

Antonio Mariscal Martínez

1. Giuliano AE, Kirgan DM, Guenther JM, Morton DL. Lymphatic mapping and sentinel lymphadenectomy for breast cancer. *Ann Surg* 1994; 220: 391-401.
2. Turner RR, Ollila DW, Krasne DL, Giuliano AE. Histopathologic validation of the sentinel lymph node hypothesis for breast cancer. *Ann Surg* 1997; 226: 271-8.
3. Albertini JJ, Lyman GH, Cox C, Yeatman T, Balducci L, Ku N, et al. Lymphatic mapping and sentinel node biopsy in the patient with breast cancer. *JAMA* 1996; 276: 1818-22.
4. Cox CE, Pendas S, Cox JM, Joseph E, Shons AR, Yeatman T, et al: Guidelines for sentinel node biopsy and lymphatic mapping of patients with breast cancer. *Ann Surg* 1998; 227: 645-53.
5. Krag D, Weaver D, Ashikaga T, Moffat F, Klimberg VS, Shriver C, et al. The sentinel node in breast cancer a multicenter validation study. *N Engl J Med* 1998; 339: 941-6.
6. Veronesi U, Paganelli G, Viale G, Galimberti V, Luini A, Zurrada S, et al. Sentinel lymph node biopsy and axillary dissection in breast cancer: results in a large series. *J Natl Cancer Inst* 1999; 91: 368-373.
7. Viale G, Bosari S, Mazzarol G, Galimberti V, Luini A, Veronesi P, et al. Intraoperative examination of axillary sentinel lymph nodes in breast carcinoma patients. *Cancer* 1999; 85: 2433-8.
8. Hill ADK, Tran KN, Akhurst T, Yeung H, Yeh SDJ, Rosen PP, et al. Lessons learned from 500 cases of lymphatic mapping for breast cancer. *Ann Surg* 1999; 229:528-535.
9. Valdes-Olmos RA, Jansen L, Hoefnagel CA, Nieweg OE, Muller SH, Rutgers EJ, et al. Evaluation of mammary lymphoscintigraphy by a single intratumoral injection for sentinel node identification. *J Nucl Med*. 2000; 41: 1500-6.
10. Veronesi U, Paganelli G, Viale G, Luini A, Zurrada S, Galimberti V, et al. A randomized comparison of sentinel-node biopsy with routine axillary dissection in breast cancer. *N Engl J Med*. 2003; 349:546-53.
11. Lyman GH, Giuliano AE, Somerfield MR, Benson AB, Bodurka DC, Burstein HJ et al. American Society of Clinical Oncology guideline recommendations for sentinel lymph node biopsy in early-stage breast cancer. *J Clin Oncol* 2005; 23: 7703-20.

12. van Rijk MC, Tanis PJ, Nieweg OE, Olmos RA, Rutgers EJ, Hoefnagel CA, Et al. Clinical implications of sentinel nodes outside the axilla and internal mammary chain in patients with breast cancer. *J Surg Oncol* 2006; 94: 281-6.
13. Kim T, Giuliano AE, Lyman GH. Lymphatic mapping and sentinel lymph node biopsy in early-stage breast carcinoma: a metaanalysis. *Cancer*. 2006; 106: 4-16.
14. Xing Y, Foy M, Cox DD, Kuerer HM, Hunt KK, Cormier JN. Meta-analysis of sentinel lymph node biopsy after preoperative chemotherapy in patients with breast cancer. *Br J Surg*. 2006; 93: 539-546.
15. van Rijk MC, Nieweg OE, Rutgers EJ, Oldenburg HS, Olmos RV, Hoefnagel CA, Kroon BB. Sentinel node biopsy before neoadjuvant chemotherapy spares breast cancer patients axillary lymph node dissection. *Ann Surg Oncol* 2006; 13: 475-9.
- 16.
17. Piñero A, Giménez J, Merck B, Vázquez C. Reunión de Consenso sobre la biopsia selectiva del ganglio centinela en el cáncer de mama. *Sociedad Española de Senología y Patología Mamaria. Rev Esp Med Nucl*. 2007;26:176-80.
18. Intra M, Rotmeasz N, Veronesi P, Colleoni M, Iodice S, Paganelli G Sentinel node biopsy is not a standard procedure in ductal carcinoma in situ of the breast: the experience on 854 patients in 10 years. *Ann Surg* 2008; 247: 315-9.
19. Mariscal A, Balliu E, Solá A, Pérez A, Fraile M, Julián JF. Comparación entre la inyección intratumoral y peritumoral del radiotrazador para la localización y biopsia del ganglio centinela en cáncer de mama no palpable. *Radiología* 2008; 50: 309-16.
20. Papa MZ, Zippel D, Kaufman B, Shimon-Paluch S, Yosepovich A, Oberman B, et al. Timing of sentinel lymph node in patients receiving neoadjuvant chemotherapy for breast cancer. *J Surg Oncol*. 2008; 98:403-6.
21. Zavagno G, De Salvo GL, Scalco G, Bozza F, Barutta L, Del Bianco P, et al. GIVOM Trialists. A Randomized clinical trial on sentinel lymph node biopsy versus axillary lymph node dissection in breast cancer: results of the Sentinella/GIVOM trial. *Ann Surg*. 2008; 247:207-13.
22. Solá M, Fraile M, Mariscal A, Julián FJ, Gubern JM, Culell P et al. Estudio comparativo de la biopsia del ganglio centinela en pacientes con carcinoma de mama multifocal vs pacientes con carcinoma unifocal. *Radiología* 2009; 51: 140-7.

23. Hansen NM, Grube B, Ye X, Turner RR, Brenner RJ, Sim MS, Giuliano AE. Impact of micrometastases in the sentinel node of patients with invasive breast cancer. *J Clin Oncol.* 2009;27:4679-84.

24. Yegiyants S, Romero LM, Haigh PI, DiFronzo LA. Completion axillary lymph node dissection not required for regional control in patients with breast cancer who have micrometastases in a sentinel node. *Arch Surg.* 2010;145:564-9.

25. Veronesi U, Viale G, Paganelli G, Zurrada S, Luini A, Galimberti V, et al. Sentinel lymph node biopsy in breast cancer: ten-year results of a randomized controlled study. *Ann Surg.* 2010; 251:595-600.